

MARIPRIST

Mifepristone (200mg) +
4 tablets Misoprostol (200mcg)

High quality, effective, proven
For abortion of early pregnancy up to 9 weeks



Your Health, Your Choice, Your Future



For early abortion of pregnancy up to 9 weeks. **Up to 98% effective when used according to instructions.**

How does it work? The first tablet causes the pregnancy to separate from the womb. The next 4 tablets cause cramps and empty the womb.

How to take Mariprist

Day 1: swallow **1 tablet with water**

Wait for **24-48 hours**

Day 2: place **4 tablets under the tongue** (sublingually) and allow to dissolve for 30 minutes. Swallow the remainder with water.

When to take Mariprist

If you are less than 9 weeks pregnant

less than 9 weeks

When not to take Mariprist

If you are more than 9 weeks pregnant

If you have severe or persistent abdominal pain

If you have an IUD in your womb

What to expect after taking Mariprist and possible side effects

Moderate bleeding with passage of clots

Shivering

Cramping

Vomiting/ diarrhoea

Warning signs: if experienced, seek medical help

These complications are rare

Heavy bleeding or no bleeding

High fever

Excess cramping /abdominal pain

Excess vomiting/diarrhoea



For more information, please scan the QR code below:
Or visit us at: www.mariprist.com

SCAN FOR MORE INFO



MARIPRIST

Mifepristone (200mg) +
4 tablets Misoprostol (200mcg)

High quality, effective, proven
For abortion of early pregnancy up to 9 weeks



Your Health, Your Choice, Your Future



For early abortion of pregnancy up to 9 weeks. **Up to 98% effective when used according to instructions.**

How does it work? The first tablet causes the pregnancy to separate from the womb. The next 4 tablets cause cramps and empty the womb.

How to take Mariprist

Day 1: swallow **1 tablet with water**

Day 2: place **4 tablets under the tongue** (sublingually) and allow to dissolve for 30 minutes. Swallow the remainder with water.

When to take Mariprist

If you are less than 9 weeks pregnant

less than 9 weeks

When not to take Mariprist

If you are more than 9 weeks pregnant

If you have severe or persistent abdominal pain

If you have an IUD in your womb

What to expect after taking Mariprist and possible side effects

Moderate bleeding with passage of clots

Shivering

Cramping

Vomiting/diarrhoea

Warning signs: if experienced, seek medical help

These complications are rare

Heavy bleeding or no bleeding

High fever

Excess cramping/abdominal pain

Excess vomiting/diarrhoea



For more information, please scan the QR code below:
Or visit us at: www.mariprist.com

SCAN FOR MORE INFO

